
Pancreatic beta-cell neogenesis by direct conversion from mature alpha-cells.

Journal: Stem Cells

Publication Year: 2010

Authors: Cheng-Ho Chung, Ergeng Hao, Ron Piran, Ehud Keinan, Fred Levine

PubMed link: 20653050

Funding Grants: Type III CIRM Stem Cell Research Training Program

Public Summary:

Because type 1 and type 2 diabetes are characterized by loss of beta-cells, beta-cell regeneration has garnered great interest as an approach to diabetes therapy. Here, we developed a new model of beta-cell regeneration, combining pancreatic duct ligation (PDL) with elimination of pre-existing beta-cells with alloxan. In this model, in which virtually all beta-cells observed are neogenic, large numbers of beta-cells were generated within 2 weeks. Strikingly, the neogenic beta-cells arose primarily from alpha-cells. alpha-cell proliferation was prominent following PDL plus alloxan, providing a large pool of precursors, but we found that beta-cells could form from alpha-cells by direct conversion with or without intervening cell division. Thus, classical asymmetric division was not a required feature of the process of alpha- to beta-cell conversion. Intermediate cells coexpressing alpha-cell- and beta-cell-specific markers appeared within the first week following PDL plus alloxan, declining gradually in number by 2 weeks as beta-cells with a mature phenotype, as defined by lack of glucagon and expression of MafA, became predominant. In summary, these data revealed a novel function of alpha-cells as beta-cell progenitors. The high efficiency and rapidity of this process make it attractive for performing the studies required to gain the mechanistic understanding of the process of alpha- to beta-cell conversion that will be required for eventual clinical translation as a therapy for diabetes.

Scientific Abstract:

Because type 1 and type 2 diabetes are characterized by loss of beta-cells, beta-cell regeneration has garnered great interest as an approach to diabetes therapy. Here, we developed a new model of beta-cell regeneration, combining pancreatic duct ligation (PDL) with elimination of pre-existing beta-cells with alloxan. In this model, in which virtually all beta-cells observed are neogenic, large numbers of beta-cells were generated within 2 weeks. Strikingly, the neogenic beta-cells arose primarily from alpha-cells. alpha-cell proliferation was prominent following PDL plus alloxan, providing a large pool of precursors, but we found that beta-cells could form from alpha-cells by direct conversion with or without intervening cell division. Thus, classical asymmetric division was not a required feature of the process of alpha- to beta-cell conversion. Intermediate cells coexpressing alpha-cell- and beta-cell-specific markers appeared within the first week following PDL plus alloxan, declining gradually in number by 2 weeks as beta-cells with a mature phenotype, as defined by lack of glucagon and expression of MafA, became predominant. In summary, these data revealed a novel function of alpha-cells as beta-cell progenitors. The high efficiency and rapidity of this process make it attractive for performing the studies required to gain the mechanistic understanding of the process of alpha- to beta-cell conversion that will be required for eventual clinical translation as a therapy for diabetes.

Source URL: <http://www.cirm.ca.gov/about-cirm/publications/pancreatic-beta-cell-neogenesis-direct-conversion-mature-alpha-cells>